USDA National Nutrient Database for Standard Reference Release 27

Statistics Report 12023, Seeds, sesame seeds, whole, dried

Report Date:October 09, 2015 17:41 EDT

Nutrient values and weights are for edible portion.

| Nutrient | Unit | Value Per100 g | Data Points S | td. Error | Min | Max | df | LB | UB | # Studies | Source | NDB Ref | Last Modified |
|-----------------------------|------|-------------------|------------------|-----------|------|------|-----|-------|-------|-----------|--|---------|------------------|
| Proximates | | | | | | | | | | | | | |
| Water | g | 4.69 | 65 | 0.180 | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Energy | kcal | 573 | | | | | | | | | Calculated or imputed | | 09/1984 |
| Energy | kJ | 2397 | | | | | | | | | Calculated or imputed | | 03/2009 |
| Protein | g | 17.73 | 61 | 0.256 | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Total lipid (fat) | g | 49.67 | 58 | 0.501 | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Ash | g | 4.45 | 59 | 0.152 | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Carbohydrate, by difference | g | 23.45 | | | | | | | | | Calculated or imputed | | 09/1984 |
| Fiber, total dietary | g | 11.8 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Sugars, total 1 | g | 0.30 | | 0.006 | 0.29 | 0.31 | 2.0 | 0.275 | 0.325 | 1 | Analytical or derived from analytical | | 02/2003 |

Minerals

| Nutrient | Unit | Value Per100 g | Data Points | Std. Error | Min | Max | df | LB | UB | # Studies | Source | NDB Ref | Last Modified |
|------------------------|------|-------------------|----------------|------------|------|------|-----|--------|--------|-----------|--|---------|------------------|
| Calcium, Ca | mg | 975 | | 102.528 | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Iron, Fe | mg | 14.55 | | 1.587 | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Magnesium, Mg | mg | 351 | | 23.961 | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Phosphorus, P | mg | 629 | | 23.830 | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Potassium, K | mg | 468 | | 34.612 | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Sodium, Na | mg | 11 | | 6.384 | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Zinc, Zn | mg | 7.75 | | 0.536 | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Copper, Cu | mg | 4.082 | | 1.277 | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Manganese, Mn | mg | 2.460 | | 0.723 | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Selenium, Se 2 3 4 5 6 | μg | 34.4 | | 6.162 | 15.6 | 97.5 | 2.0 | 11.659 | 57.127 | 5 | Analytical or derived from analytical | | 03/2009 |

Vitamins

| Nutrient | Unit | Value Per100 g | Data Points S | td. Error | Min | Max | df | LB | UB | # Studies | Source | NDB Ref | Last Modified |
|--------------------------------|------|-------------------|------------------|-----------|-----|-----|----|----|----|-----------|--|---------|------------------|
| Vitamin C, total ascorbic acid | mg | 0.0 | | 0.000 | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Thiamin | mg | 0.791 | | 0.084 | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Riboflavin | mg | 0.247 | | 0.034 | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Niacin | mg | 4.515 | | 0.227 | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Pantothenic acid | mg | 0.050 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Vitamin B-6 | mg | 0.790 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Folate, total | μg | 97 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Folic acid | μg | 0 | | | | | | | | | Assumed zero | | 01/2001 |
| Folate, food | μg | 97 | | | | | | | | | Analytical or derived from analytical | | 03/2009 |
| Folate, DFE | μg | 97 | | | | | | | | | Calculated or imputed | | 03/2009 |
| Choline, total | mg | 25.6 | | | | | | | | | Calculated or imputed | 12201 | 12/2006 |
| Vitamin B-12 | μg | 0.00 | | | | | | | | | Assumed zero | | 09/1984 |

| Nutrient | Unit | Value Per100 g | Data Points St | d. Error | Min | Max | df | LB | UB | # Studies | Source | NDB Ref | Last Modified |
|-----------------------------------|------|-------------------|-------------------|----------|-----|-----|----|----|----|-----------|--|---------|------------------|
| Vitamin B-12, added | μg | 0.00 | | | | | | | | | Assumed zero | | 09/2004 |
| Vitamin A, RAE | μg | 0 | | | | | | | | | Calculated or imputed | | 03/2009 |
| Retinol | μg | 0 | | | | | | | | | Assumed zero | | 06/2002 |
| Carotene, beta | μg | 5 | | | | | | | | | Calculated or imputed | | 02/2003 |
| Carotene, alpha | μg | 0 | | | | | | | | | Assumed zero | | 02/2003 |
| Cryptoxanthin, beta | μg | 0 | | | | | | | | | Assumed zero | | 02/2003 |
| Vitamin A, IU | IU | 9 | | 6.094 | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Lycopene | μg | 0 | | | | | | | | | Assumed zero | | 02/2003 |
| Lutein + zeaxanthin | μg | 0 | | | | | | | | | Assumed zero | | 02/2003 |
| Vitamin E (alpha-tocopherol) | mg | 0.25 | | | | | | | | | Calculated or imputed | | 02/2003 |
| Vitamin E, added | mg | 0.00 | | | | | | | | | Assumed zero | | 09/2004 |
| Vitamin D (D2 + D3) | μg | 0.0 | | | | | | | | | Assumed zero | | 03/2009 |
| Vitamin D | IU | 0 | | | | | | | | | Assumed zero | | 03/2009 |
| Vitamin K (phylloquinone) Lipids | μg | 0.0 | | | | | | | | | Calculated or imputed | 12201 | 02/2003 |
| Fatty acids, total saturated | g | 6.957 | | | - | | | | | | Analytical or derived from analytical | | 09/1984 |
| 4:0 | g | 0.000 | | | | | | | | | Analytical or derived from analytical | | 02/1995 |

| Nutrient | Unit | Value Per100 g | Data Points Std. E | Error | Min | Max | df | LB | UB | # Studies | Source | NDB Ref | Last Modified |
|------------------------------------|------|-------------------|-----------------------|-------|-----|-----|----|----|----|-----------|--|---------|------------------|
| 6:0 | g | 0.000 | | | | | | | | | Analytical or derived from analytical | | 02/1995 |
| 8:0 | g | 0.000 | | | | | | | | | Analytical or derived from analytical | | 02/1995 |
| 10:0 | g | 0.000 | | | | | | | | | Analytical or derived from analytical | | 02/1995 |
| 12:0 | g | 0.000 | | | | | | | | | Analytical or derived from analytical | | 02/1995 |
| 14:0 | g | 0.124 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| 16:0 | g | 4.441 | 761 | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| 18:0 | g | 2.090 | 761 | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Fatty acids, total monounsaturated | g | 18.759 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| 16:1 undifferentiated | g | 0.149 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| 18:1 undifferentiated | g | 18.521 | 761 | | | | | | | | Analytical or derived from analytical | | 09/1984 |

| Nutrient | Unit | Value Per100 g | Data Points Std. | Error | Min | Max | df | LB | UB | # Studies | Source | NDB Ref | Last Modified |
|------------------------------------|------|-------------------|---------------------|-------|-----|-----|----|----|----|-----------|--|---------|------------------|
| 20:1 | g | 0.070 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| 22:1 undifferentiated | g | 0.000 | | | | | | | | | Analytical or derived from analytical | | 02/1995 |
| Fatty acids, total polyunsaturated | g | 21.773 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| 18:2 undifferentiated | g | 21.375 | 761 | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| 18:3 undifferentiated | g | 0.376 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| 18:4 | g | 0.000 | | | | | | | | | Analytical or derived from analytical | | 02/1995 |
| 20:4 undifferentiated | g | 0.000 | | | | | | | | | Analytical or derived from analytical | | 02/1995 |
| 20:5 n-3 (EPA) | g | 0.000 | | | | | | | | | Analytical or derived from analytical | | 02/1995 |
| 22:5 n-3 (DPA) | g | 0.000 | | | | | | | | | Analytical or derived from analytical | | 02/1995 |
| 22:6 n-3 (DHA) | g | 0.000 | | | | | | | | | Analytical or derived from analytical | | 02/1995 |

| Nutrient | Unit | Value Per100 g | Data Points Std | . Error | Min | Max | df | LB | UB | # Studies | Source | NDB Ref | Last Modified |
|---------------------------|------|-------------------|--------------------|---------|-----|-----|----|----|----|-----------|--|---------|------------------|
| Cholesterol | mg | 0 | | | | | | | | | Assumed zero | | 09/1984 |
| Phytosterols Amino Acids | mg | 714 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Tryptophan | g | 0.388 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Threonine | g | 0.736 | | | | | | | | | Analytical or derived from analytical | - | 09/1984 |
| Isoleucine | g | 0.763 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Leucine | g | 1.358 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Lysine | g | 0.569 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Methionine | g | 0.586 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Cystine | g | 0.358 | - | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Phenylalanine | g | 0.940 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |

| Nutrient | Unit | Value Per100 g | Data Points Std. En | rror | Min | Max | df | LB | UB | # Studies | Source | NDB Ref | Last Modified |
|---------------|----------------|-------------------|------------------------|------|-----|-----|----|----|----|-----------|--|---------|------------------|
| Tyrosine | σg | 0.743 | - | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Valine | g | 0.990 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Arginine | g | 2.630 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Histidine | g | 0.522 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Alanine | g | 0.927 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Aspartic acid | g | 1.646 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Glutamic acid | g | 3.955 | - | | | - | | | | | Analytical or derived from analytical | | 09/1984 |
| Glycine | g | 1.215 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Proline | o _D | 0.810 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |
| Serine | g | 0.967 | | | | | | | | | Analytical or derived from analytical | | 09/1984 |

Other

USDA National Nutrient Database for Standard Reference Release 27 Statistics Report October 09, 2015 17:41 EDT Page 9 of 9

| Nutrient | Unit | Value Per100 g | Data Points | Std. Error | Min | Max | df | LB | UB | # Studies | Source | NDB Ref | Last Modified |
|-----------------------|------|-------------------|----------------|------------|------|------|----|----|----|-----------|--------------|---------|------------------|
| Alcohol, ethyl | g | 0.0 | | | | | | | | | Assumed zero | | 04/1985 |
| Caffeine | mg | 0 | | | | | | | | | Assumed zero | | 02/2003 |
| Theobromine | mg | 0 | | | | | | | | | Assumed zero | | 02/2003 |
| Flavonoids | | | | | | | | | | | | | |
| Isoflavones | | | | | | | | | | | | | |
| Daidzein 7 8 | mg | 0.0 | | | 0 | 0 | | | | | | | |
| Genistein 7 8 | mg | 0.0 | | | 0 | 0 | | | | | | | |
| Glycitein 8 | mg | 0.0 | | | 0 | 0 | | - | | | | | |
| Total isoflavones 7 8 | mg | 0.0 | | | 0.01 | 0.01 | | | | | | | |
| Formononetin | mg | 0.0 | | | 0 | 0 | | - | | | | | |
| Coumestrol | mg | 0.0 | | | 0 | 0 | | | | | | | |

Sources of Data

¹Nutrient Data Laboratory, ARS, USDA Variability of the sugar content of foods, 1989 Beltsville MD

²E.C. Pappa, A.C. Pappas, P.F. Surai Selenium content in selected foods from the Greek market and estimation of the daily intake, 2006 Science of the Total Environment 372 pp.100-108

³P.P. Sirichakwal, P. Puwastien, J. Polngam, R. Kongkachuichai Selenium content of Thai foods, 2005 J Food Comp Anal 18 pp.47-59

⁴R. Abdulah, K. Miyazaki, M. Nakazawa, H. Koyama Low contribution of rice and vegetables to the daily intake of selenium in Japan, 2005 International Journal of Food Science and Nutrition 56 7 pp.463-471

⁵A. Askar, H.J. Bielig Selenium content of food consumed by Egyptians, 1983 Food Chemistry 10 pp.231-234

⁶Nutrient Data Laboratory, ARS, USDA National Food and Nutrient Analysis Program Wave 6m, 2002 Beltsville MD

⁷Liggins, J., Bluck, L. J. C., Runswick, S., Atkinson, C., Coward, W. A., Bingham, S. A. Daidzein and genistein content of fruits and nuts., 2000 J. Nutr. Biochem. 11 pp.326-331

⁸Thompson, L. U., Boucher, B. A., Liu, Z., Cotterchio, M., and Kreiger, N. Phytoestrogen content of foods consumed in Canada, including isoflavones, lignans, and coumestan., 2006 Nutr. Cancer 54 pp.184-201